

DISCRIMINATION OF MALAYSIAN SWIFLET'S NEST AND IT'S POTENTIAL ADULTERANTS USING HANDHELD PORTABLE FTIR SPECTROSCOPIC SYSTEM

G. S. L. Ketty, F. H. Roosnoor, I. Faridah, and I. Norakmar

Veterinary Public Health Laboratory, Department of Veterinary Services, Jln Nilai-Banting, Bandar Baru Salak Tinggi, 43900, Sepang, Selangor

INTRODUCTION

Edible bird nest (EBN) is woven from gelatinous strands of Swiftlet's saliva mixed with minor feathers during their breeding season (1). The Chinese has been enjoying this cuisine for thousands of years back since Tang (618-907 AD) and Sung (960-1279 AD) dynasties. EBN can be sold at very high price due to its high nutritional and medicinal values. Besides, it is also considered as the world most expensive animal product (2). These reasons led to rampant adulteration of the expensive pure EBN. Various adulterants can be added into the pure EBN which includes Tremella fungus, karaya gum, red seaweed, fried pork skin, egg white, gelatin, vermicelli rice, rice, cellophane noodles, starch, milk, and jelly. Other adulteration methods include staining, bleaching and incorporating cheaper EBN into the more expensive ones (3; 4; 5).

A variety of authentication techniques had been established (4) and all have their own advantages and disadvantages. The empirical method is rather simple compared to chemical method that requires extraction of the EBN prior to analysis and mostly not environmental friendly.

Handheld Portable Fourier Transform Infrared (FTIR) system can be used in the laboratory and in the field in the same manner with no sample preparation needed (6). This system is versatile, robust and come with interchangeable sampling interfaces; diffuse, grazing angle, specular reflection or spherical attenuated total reflectance (ATR) that make it a highly useful handheld mid-IR spectrometer that suits with any type of samples. A resulting spectrum represents molecular absorption and transmission, creating a molecular fingerprint of the EBN. Like any fingerprint, no two unique molecular structures produce the same infrared spectrum.

This study was carried out to discriminate the EBN and its' potential adulterants through the EBN fingerprint using the handheld portable FTIR system.

MATERIALS AND METHODS



RESULTS AND DISCUSSION




Figure 1: Fingerprint of pure EBN and its' potential adulterants




Figure 2: Fingerprint of pure EBN and adulterated EBN

- The spectra showed intense peaks and several spectra visibly overlapped. The most characteristic FTIR absorption bands of pure EBN were found at about 1640 cm^{-1} to 1320 cm^{-1} representing the secondary protein which is the amide (CO-NH_2), mono substitution amide (CO-NH-R) and di substitution amide (CO-NR_2). Another major band observed was carbohydrate at 1090 cm^{-1} until 900 cm^{-1} .
- The adulterants and adulterated EBN fingerprint spectra did not have all the amides while the carbohydrate peak is higher compared to pure EBN fingerprint which indicated that the carbohydrate concentration was higher (7). However, the seaweed and adulterated EBN with seaweed had lesser carbohydrate bond. These correlates with high crude protein level in pure EBN and high carbohydrate level in the adulterants and adulterated EBN except for seaweed (8).
- An EBN purity library was created by inserting all the pure EBN, adulterants and adulterated EBN fingerprint spectra. The library was tested using other pure EBN and similarities were found to be more than 90%.



Spectral	• 4000-650 cm^{-1}
No. of Scan	• 64 times
Resolution	• 4 cm^{-1}
Sampling Interface	• diffuse reflectance
Software	• MicroLab • Spekwin32 1.71.6.1

MicroLab PC software (Agilent) Setting



Scanning
↓
Spectrum Interpretation using Structure Correlation Charts



EBN Purity Library using MicroLab

REFERENCES

- (1) Koon, L. C. (2000). Features – Bird's Nest Soup – Market demand for this expensive gastronomic delicacy threatens the aptly named edible-nest Swiftlets with extinction in the east. *Wildlife Conservation* 103(1): 30-35.
- (2) Koon, L. C., & Cranbrook, Earl of (2002). Swiftlets of Borneo – Builders of edible nests (pp.1-171). Sabah, Malaysia: Natural History Publication (Borneo) Sdn. Bhd..
- (3) Joe, S. (2012). Fast, effective evaluation of edible bird nests using handheld Agilent 4100 ExoScan FTIR. Application note from Agilent Technologies, Inc.
- (4) Ma, F. C., & Liu, D. C. (2012). Extraction and determination of hormones in the edible bird's nest. *Asian Journal of Chemistry*, 24(1), 117-120
- (5) Marcone, F. M. (2005). Characterization of the edible bird's nest the "Caviar of the East". *Food Research International*, 38 (2005) 1125 – 1134.
- (6) Agilent Technologies. 4100 ExoScan Series FTIR (handheld). <http://www.chem.agilent.com/en-US/products-services/Instruments-Systems/Molecular-Spectroscopy/4100-ExoScan-Series-FTIR-%28handheld%29/Pages/default.aspx>
- (7) Ketty, G. S. L., Roosnoor, F. H., and Faridah, I. (2013). Rapid and Non-Destructive Screening Method for Pure Malaysian Swiftlet's Nest and It Potential Adulterants using Handheld FTIR. 25th Veterinary Association Malaysia (VAM) 2013, Lotus Desaru Beach Resort, Desaru, Johor, Malaysia.
- (8) Marcone, F. M. (2011). *Origins and Compositional Analysis of Novel Foods: Kopi Luwak Coffee and Bird's Nest Soup*. Elsevier B. V.

CONCLUSION

- Based on the results, the FTIR spectroscopy with the purity library created demonstrated that this technique is nondestructive, effective, simple and reliable. Unlike other common techniques, handheld portable FTIR-diffuse spectroscopy can be used in the field as it does not require laborious sample preparation procedures and the results can be obtained by simple interpretation from the library.

Researcher information



Dr. Ketty Goh Siew Lin
Veterinary Public Health Laboratory,
Jalan Nilai-Banting, B.B. Salak Tinggi,
43900 Sepang, Selangor
Tel: 03-87068681 Fax: 03-87068675
Email: ketty@dvs.gov.my

ACKNOWLEDGEMENTS

The authors would like to thanks Director General of Veterinary Services, Director of BDKK and Deputy Director of VPHL for the permission to publish the findings of this study at EBNIC 2014. The authors wish to acknowledge the technical assistance provided by the laboratory staff of Additive Unit, VPHL. Special thanks to Exotic Animal Section, DVS for providing the FTIR systems and raw EBN samples.